

GROUP 1 HERBICIDE

PULL HERE TO OPEN ►

 **Fusion**®

**syngenta**®

## Herbicide

Postemergence Herbicide for Control of Perennial and Annual Grass Weeds

**Active Ingredients:**

Fluazifop-P-butyl	
Butyl( <i>R</i> )-2-[4-[[5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]propanoate	24.15%
Fenoxaprop-P-ethyl	
(+)-ethyl-2-[4-[6-(chloro-2-benzoxazolyl)oxy]phenoxy]propanoate	6.76%
Other Ingredients*	69.09%
<b>Total:</b>	<b>100.00%</b>

Contains 2 lbs. (+) isomer (fluazifop-P-butyl) and 0.56 lbs. fenoxaprop-P-ethyl active ingredient per gallon.

\*Contains aromatic petroleum distillates.

**KEEP OUT OF REACH OF CHILDREN.**

### WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See directions for use inside booklet.

**EPA Reg. No. 100-1059 EPA Est. No. 11773-IA-01**

**Product of United Kingdom  
Formulated in the USA**

**SCP 1059A-L1D 1210  
334497**

**2.5 gallons**  
Net Contents

TM

<b>FIRST AID</b>	
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Immediately call a poison control center or doctor.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give any liquid to the person.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>NOTE TO PHYSICIAN</b>	
Contains petroleum distillate – vomiting may cause aspiration pneumonia.	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>HOT LINE NUMBER</b>	
For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident) Call <b>1-800-888-8372</b>	

## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

#### WARNING/AVISO

WARNING. Causes skin irritation. Harmful if absorbed through skin or swallowed. Causes moderate eye irritation. Do not get on skin or on clothing. Avoid contact with eyes. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

#### Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant shoes plus socks, and
- Chemical-resistant gloves, such as barrier laminate or Viton®

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

*continued...*

## PRECAUTIONARY STATEMENTS (*continued*)

### User Safety Recommendations

#### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### Environmental Hazards

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Do not apply when weather conditions favor drift from target area.

### Ground Water Advisory

Fluazifop-P-butyl is known to leach through soil into ground water under certain conditions as a result of label use. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water.

This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of fluazifop-p-butyl from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

### Physical and Chemical Hazards

**Combustible.** Do not use or store near heat or open flame.

## CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold Syngenta and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, In no event shall Syngenta be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or Viton
- Shoes plus socks

## PRODUCT INFORMATION

Read all label directions before using.

Fusion Herbicide is a selective postemergence herbicide for control of annual and perennial grass weeds in soybeans, and cotton. Fusion Herbicide provides effective control of grass weeds in conventional tillage, minimum tillage, and no-till plantings. Fusion Herbicide may be applied for control of emerged grass weeds before, during or after planting or after harvest of soybeans or cotton. Fusion Herbicide does not control broadleaf weeds or sedges (nutgrass).

Fusion Herbicide is a systemic herbicide which moves from the treated foliage into the shoots, roots, rhizomes, stolons, and growing points (meristematic regions) of treated grass weeds.

**Rainfastness** – Since Fusion Herbicide is rapidly absorbed by the grass foliage, rain or irrigation occurring one hour or more after application will not affect the activity of Fusion Herbicide. When tank mixing with broadleaf herbicides, observe the rainfast statement of the most restrictive label.

**Control Symptoms** – Growth of treated grass weeds stops soon after application. Symptoms include loss of vigor, yellowing and/or reddening, and eventual death of the treated grass plant. Symptoms are generally observed within one week after treatment, depending on grass weed species and environmental conditions.

#### **INFORMATION ON WEED RESISTANCE**

Naturally occurring biotypes of certain grass species with resistance to this herbicide and related products (same mode of action) are known to exist. Selection of resistant biotypes, through repeated use of these herbicides, may result in control failures.

If poor performance cannot be attributed to adverse weather conditions or improper application methods, a resistant biotype may be present. In such a case, additional treatments with this herbicide or related products are not recommended. Consult your local company representative or agricultural advisor for assistance.

#### **APPLICATION DIRECTIONS**

Thorough coverage of all grass weed foliage is important for good activity. Optimum weed control is achieved when young actively growing grass weeds are treated that are not under stress from low soil moisture, extreme temperatures, low soil fertility, mechanical, or chemical injury.

**Spray Additives** - Only crop oil concentrate, nonionic surfactants and other adjuvants cleared for use on growing crops may be used in spray mixture. Under dry conditions, crop oil concentrate is the preferred adjuvant.

##### **Always Add One of the Following:**

- **Crop Oil Concentrate (COC)** – Add a non-phytotoxic crop oil concentrate or a once-refined vegetable oil concentrate containing 15-20% approved emulsifier, at 0.5-1% v/v (0.5-1 gal./100 gals.) in the finished spray volume for ground applications. For aerial applications, add 1 pint of crop oil concentrate per acre.
- **Nonionic Surfactant (NIS)** – Add nonionic surfactant containing at least 75% surface-active agent, at 0.25-0.5% v/v (1-2 qts./100 gals.) in the finished spray volume for ground application. For aerial application, add 1 pint of surfactant per acre.
- **Other Adjuvants** - Adjuvants other than COC or NIS may be used providing the product meets the following criteria:
  1. Contains only EPA exempt ingredients.
  2. Is nonphytotoxic to the target crop.
  3. Is compatible in mixture. (May be established through a jar test.)
  4. Is supported locally for use with Fusion on the target crop through proven field trials and through university and extension recommendations.

Always refer to the product label and follow directions concerning rates, target crops, environmental effect such as drought or weed stress, and use in tank mix with other labeled pesticides.

#### **DIRECTIONS FOR GROUND APPLICATION**

**Nozzle Selection** - The use of flat fan nozzles will result in the most effective application of Fusion Herbicide. Flood nozzles are generally not as good as flat fans since they produce large uneven droplets. The use of nozzles other than flat fans may result in reduced grass control due to inadequate coverage. Do not apply Fusion Herbicide with recirculating sprayers, rope-wick applicators, controlled droplet applicators (CDA) or any similar devices.

**Spray Volume and Pressure** - Use sufficient spray volume and pressure to ensure complete coverage of the target grasses. Apply in 5-40 gallons per acre of spray mixture with spray pressures of 30-60 psi at the nozzle tip. When grass foliage is dense, use 60 psi and a minimum of 20 gallons per acre to ensure coverage of weed foliage.

**Band Applications** - Use a minimum of two nozzles, one directed to each side of the planted row. Application with a single nozzle directed over the top of the row is not recommended. Cultivation of untreated areas may be needed following band applications.

When making band applications and cultivating in the same operation, position nozzles in advance of the cultivation device. This will reduce dust in the spray area. Dust can intercept the spray, reducing weed coverage resulting in less than adequate weed control.

Calculate the amount of herbicide and water volume needed for band treatment by the following formulas:

$$\frac{\text{Band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{Band herbicide rate per acre}$$

$$\frac{\text{Band width in inches}}{\text{row width in inches}} \times \text{broadcast volume per acre} = \text{Band water volume per acre}$$

Band applications to perennial grasses are not recommended, as reinfestation of the treated band from the untreated middle may result.

**Spot Treatments** – Mix Fusion Herbicide and crop oil concentrate or nonionic surfactant with water according to the amounts shown below. Spray to obtain thorough coverage, but do not spray to runoff. Re-treat if necessary.

**Table 1. Spot Spray Mixing Directions**

To Make This Spray Volume	Add These Amounts		
	Fusion	Crop Oil Concentrate	OR Nonionic Surfactant
1 gallon	$\frac{3}{4}$ fl. oz. (1 $\frac{1}{2}$ tbsp.)	1 $\frac{1}{2}$ fl. oz.	$\frac{1}{2}$ fl. oz.
10 gallons	6.5 fl. oz.	13 fl. oz.	3 fl. oz.
25 gallons	1 pt.	1 qt.	$\frac{1}{2}$ pt.
50 gallons	1 qt.	2 qts.	1 pt.

**Chemigation** - Do not apply Fusion Herbicide through any type of irrigation system.

#### **DIRECTIONS FOR AERIAL APPLICATION**

Use sufficient spray volume to ensure complete coverage of target grasses. A minimum of 5 gallons per acre should be applied. When grass foliage is dense, use a minimum of 10 gallons per acre to ensure coverage of weed foliage.

**Application Timing** - Best control of susceptible grasses is obtained when Fusion Herbicide is applied to actively growing grasses before they exceed the listed growth stages shown on this label. Refer to Tables 2, 3, 4 and 5 for specific directions on use rates and weed growth stages.

**Cultivation** - Cultivation of treated grasses is not recommended within 7 days prior to or within 7 days after application of Fusion Herbicide as weeds may be put under stress resulting in reduced weed control. Timely cultivation 2-3 weeks after applying Fusion Herbicide may assist weed control.

#### **Spray Drift Management**

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed  $\frac{3}{4}$  the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

#### **Importance of Droplet Size**

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversion section of this label).

#### **Controlling Droplet Size**

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released backwards, parallel to the airstream will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - For some use patterns, reducing the effective boom length to less than  $\frac{3}{4}$  of the wingspan or rotor length may further reduce drift without reducing swath width.
- **Application** - Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

#### **Swath Adjustment**

When applications are made with a cross-wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

#### **Wind**

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

#### **Temperature and Humidity**

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

#### **Temperature Inversions**

Applications should not occur during a temperature inversion, because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a connected cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

### **Sensitive Areas**

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

### **PRODUCT USE PRECAUTIONS**

- Apply to actively growing grasses. Avoid application to grasses which are stressed due to moisture, temperature, low soil fertility, mechanical or chemical injury.
- Apply at the directed rate to grasses at the listed growth stages as outlined in Tables 2, 3, 4 and 5.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment. Use the highest recommended rate for grasses in that population.
- Where irrigation is used as part of normal cropping practice, best results are usually obtained when Fusion Herbicide is applied within 7 days after irrigation.
- Best perennial grass control can be obtained if rhizomes or stolons are cut up by preplant tillage practices (disking, plowing, etc.) to stimulate maximum emergence of grass shoots.
- Do not plant rotational grass crops such as corn, sorghum, and cereals within 60 days of last application of Fusion Herbicide.
- Avoid drift to all other crops and nontarget areas. Grass crops are highly susceptible to Fusion Herbicide. Avoid application when wind velocity exceeds 15 mph. Do not make ground or aerial applications during temperature inversions.
- Tank mixes of Fusion Herbicide with other pesticides, liquid fertilizers or other additives except as specified on this label or on approved Syngenta supplemental labels may result in tank mix incompatibility, unsatisfactory performance and/or crop injury.
- Thoroughly clean spray tank with water and a commercial tank cleaner before and after each use.
- Do not apply Fusion Herbicide if rainfall is expected within 1 hour.

### **SPECIFIC RESTRICTIONS FOR SOYBEANS AND COTTON**

- Do not apply a total of more than 24 oz. (0.375 lbs. ai/A) of Fusion Herbicide per acre per season to soybeans and cotton.
- Do not apply to cotton after boll set.
- Do not harvest cotton within 90 days of Fusion application.
- Make the last Fusion Herbicide application to soybeans before bloom.
- Do not graze or harvest for forage or hay.

## **DIRECTIONS FOR ANNUAL AND PERENNIAL GRASS WEED CONTROL IN SOYBEANS AND COTTON – REGION A**

### **SOYBEANS**

Fusion Herbicide may be applied in the following states: Alabama, Arkansas, Connecticut, Delaware, Florida, Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Nebraska, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma (east of Interstate 35), Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas (east of Interstate 35), Vermont, Virginia, West Virginia, and Wisconsin.

### **COTTON**

Fusion Herbicide may be applied in the following states: Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, Missouri, North Carolina, Oklahoma (east of Interstate 35), South Carolina, Tennessee, Texas (east of Interstate 35), and Virginia.



Table 2. Fusion Herbicide Annual Grass Control Use Rate Directions in Soybeans and Cotton - Region A

Annual Grass Species <sup>1</sup>	Ht. (In.)	Special Fusion Rate When Used Alone <sup>2</sup> (fl. oz./A)	Fusion Rate When Used Alone (fl. oz./A)	Fusion Rate When Tank Mixed <sup>2</sup> (fl. oz./A)
Barnyardgrass	2-4	6	8	8-12
Broadleaf signalgrass	2-4	-	8-10	10-12
Crabgrass	1-4	-	8	8-12
Downy Brome	2-6	-	6	6-10
Fall Panicum	2-6	-	8	8-12
Field Sandbur	2-4	-	8	8-12
Foxtails				
Giant foxtail	2-8	6	7	7-12
Green foxtail	2-4	6	8	8-12
Yellow foxtail	2-4	6	8	8-12
Goosegrass	2-4	-	8	8-12
Italian Ryegrass	2-4	-	8	8-12
Itchgrass	4-24	-	6	6-10
Johnsongrass, seedling	2-8	-	6	6-10
Junglerice	2-3	-	8	8-12
Red rice	0.5-3	-	10-12	10-12
Shattercane	6-12	-	6	6-10
Sorghum almum	6-12	-	6	6-10
Southern sandbur	2-6	-	8	8-12
Texas Panicum	2-8	-	8	8-12
Volunteer Cereals				
V. Barley	2-6	-	8	8-12
V. Corn <sup>3,4</sup>	12-24	4	6	6-10
V. Milo	6-12	-	6	6-10
V. Oats	2-6	-	8	8-12
V. Rye	2-6	-	8	8-12
V. Wheat	2-6	-	8	8-12
Wild Oats	2-6	-	8	8-12
Wild Proso Millet	4-8	-	6	6-10
Witchgrass	2-4	-	8	8-12
Woolly Cupgrass	2-4	-	8	8-12

<sup>1</sup>Retreatment at the listed rate may be needed to control later germinating grasses or if regrowth occurs.

<sup>2</sup>The lowest rate of Fusion Herbicide listed above may be used under the following conditions:

- Application under favorable soil moisture and humidity conditions, normally within a few days after rainfall or irrigation. Avoid extreme air temperatures.
- Application at earliest growth stages indicated on rate tables.
- Application in highly competitive crop stands such as narrow row or drilled soybeans, or where cultivation is planned.
- Application to light or moderate weed densities.
- Application with 1% v/v crop oil concentrate only.

If the conditions above do not exist, use the high rate for the species indicated.

<sup>3</sup>Fusion will provide suppression of Sethoxydim resistant volunteer corn.

<sup>4</sup>Includes control of Roundup Ready® (glyphosate-tolerant), LibertyLink® and IMI-Corn® volunteer corn.

**Table 3. Fusion Herbicide Perennial Grass Control Use Rate Directions In Soybeans and Cotton – Region A**

Perennial Grass Species	Application	Ht. (In.)	Rate (fl. oz./A)
Bermudagrass <sup>1</sup>	1st	4-8 (Runner)	12
	2nd	4-8	8
Quackgrass <sup>2</sup>	1st	6-10	12
	2nd	10	8
Rhizome johnsongrass <sup>3</sup>	1st	8-18	10-12
	2nd	6-12	8
Wirestem muhly <sup>4</sup>	1st/2nd	4-12	10-12

<sup>1</sup>Make second application to bermudagrass if regrowth occurs (usually about 4 weeks after first application). Bermudagrass control may be improved by directing the spray beneath the crop canopy. To improve coverage, make applications in a minimum of 15 gallons per acre.

<sup>2</sup>Make second application 2-3 weeks after the first, but before the quackgrass exceeds 10 inches in height. Always use 1% v/v crop oil concentrate.

<sup>3</sup>Make first application before the boot stage. In eastern Oklahoma, the Brazos Bottoms, the Blacklands, Coastal Bend and Rio Grande areas of eastern Texas, make the first application at 8-12 inches. If new shoots emerge or regrowth occurs, make a second application at 4-6 inches.

<sup>4</sup>Make second application if regrowth occurs.

#### **RESCUE APPLICATION IN SOYBEANS**

Fusion Herbicide applied at 8-14 fl. oz./A may be used to control giant foxtail, wild proso millet and woolly cupgrass up to 16 inches in height. Applications must be made prior to soybean bloom. Use 12-14 fl. oz./A if grasses appear stressed due to drought, unfavorable temperatures and/or low soil fertility. Do not tank mix Fusion with broadleaf herbicides for rescue applications.

**DIRECTIONS FOR ANNUAL AND PERENNIAL GRASS WEED CONTROL IN COTTON – REGION B**

New Mexico, Oklahoma (West of Interstate 35), and Texas (West of Interstate 35).

**Table 4. Fusion Herbicide Annual Grass Control Use Rate Directions – Region B**

Annual Grass Species <sup>1</sup>	Ht. (In.)	No. of Leaves Not to Exceed	Rate(s) fl. oz./A
Barnyardgrass	1-2	3	12
Canarygrass, Littleseed	2-4	4	12
Crabgrass, Large	1-2	3	12
Crabgrass, Smooth <sup>2</sup>	1-2	4	12
Johnsongrass, seedling	2-4	3	8
Junglerice	2-3	3	12
Panicums			
Fall Panicum <sup>2</sup>	2-6	6	12
Texas Panicum <sup>2</sup>	8	8	12
Rabbitfootgrass	2-4	4	12
Volunteer Cereals			
V. Barley	2-4	3	12
V. Corn	12-18	6	12
V. Milo	2-4	4	12
V. Oats	2-4	3	12
V. Wheat	2-4	3	12
Wild Oats <sup>2</sup>	2-4	4	12

<sup>1</sup>Retreatment at the listed rate may be needed to control late germinating grasses or if regrowth occurs.

<sup>2</sup>Recommended for use in Oklahoma and Texas on these species.

**Table 5. Fusion Herbicide Perennial Grass Control Use Rate Directions – Region B**

For best results, apply Fusion 3 days before to 7 days after irrigation.			
Perennial Grass Species	Application	Ht. (In.)	Rate fl. oz./A
Bermudagrass <sup>1,2</sup>	1st	4-8 (runner length)	16-24
Rhizome johnsongrass <sup>3,4</sup>	1st	12-18	10-24

<sup>1</sup>Make second application to bermudagrass if regrowth occurs. (Usually about 4 weeks after first application.) Bermudagrass control may be improved by directing the spray beneath the crop canopy. To improve coverage, make applications in a minimum of 20 gallons per acre. Do not apply more than 24 oz. in one season.

<sup>2</sup>In Oklahoma (west of Interstate 35) and Texas (west of Interstate 35) under good soil moisture and favorable growing conditions, make applications at reduced rates of 12-16 fl. oz./A.

<sup>3</sup>Make first application before the boot stage. If new shoots emerge or regrowth occurs, make a second application. (Usually about 4 weeks after the first application.) Do not apply more than 24 oz. in one season.

<sup>4</sup>In Oklahoma (west of Interstate 35) and Texas (west of Interstate 35) under good soil moisture and favorable growing conditions, make the first application to 8-18 inch johnsongrass at a reduced rate of 12 fl. oz./A. Make the second application to 6-12 inch johnsongrass at a rate of 12 fl. oz./A.

## TANK MIX AND SEQUENTIAL APPLICATIONS FOR SOYBEANS

Fusion can be used sequentially or in tank mix with one or more of the following: Basagran®, Boundary®, Canopy®\*\*, Classic®\*\*, Cobra®, FirstRate®\*\*, Flexstar®, Flexstar GT, Gramoxone Inteon®, Harmony®\*\*, Ignite®, Prefix®, Prowl®, Pursuit®\*, Reflex®, Scepter®\*\*, Storm®, Synchrony®\*\*, Roundup® brands, Touchdown® brands, Ultra Blazer®, 2,4-D (LVE)\*\*\*.

\*Fusion at 4 oz./A plus Pursuit for volunteer corn and shattercane only.

\*\*If the grass population consists mainly of yellow foxtail, barnyardgrass, woolly cupgrass, field sandbur, smooth or large crabgrass, Texas panicum, broadleaf signalgrass or red rice and conditions are less than optimum (see footnote 2, Table 2), a sequential application is recommended to provide satisfactory performance.

\*\*\*Fusion + 2,4-D (LVE) tank mix at 4-8 fl. oz./A + 0.5 lb. AE/A may be used as a preplant treatment for the control of giant foxtail and fall panicum and broadleaf weeds as specified on the 2,4-D label. Use the higher rate of Fusion on grasses greater than 2 inches. The tank mix should be used with a crop oil concentrate.

Under certain conditions, the mixture of Fusion with one or more of the above-mentioned broadleaf herbicides may cause a reduction in grass activity.

For sequential applications, allow 2 to 3 days after the application of Fusion before applying a broadleaf herbicide or mixture. In case the broadleaf herbicide or mixture is applied first, apply Fusion when the grass weeds begin to develop new leaves (generally around 7 days).

Liquid nitrogen fertilizer (28% UAN or similar) may be added to the spray mixture up to 4 gallons per 100 gallons (4% v/v). Dry nitrogen fertilizer (ammonium sulfate) may be added up to 4 lbs. product per acre. Liquid and dry nitrogen fertilizers should not be used as a substitute for crop oil concentrate or nonionic surfactant in the spray mixture.

Always read and follow the application directions, restrictions and limitations for all products whether used alone or in a tank mix. The most restrictive labeling of any product used applies in tank mixtures.

**Note:** Tank mix applications sometimes have resulted in reduced grass weed control and possible increases in crop injury as compared to the products used alone. If grass regrowth occurs following an application of the tank mix or an additional flush of grasses emerge, make a second application of Fusion Herbicide to actively growing grass weeds, as per the label recommendations. When perennial grasses are the predominant grass to be controlled, a sequential application is recommended. Follow the directions for sequential applications of Fusion Herbicide and the appropriate broadleaf herbicide.

### Glyphosate-Tolerant Soybeans (including Roundup Ready Soybeans) Tank Mix

Fusion Herbicide can be tank mixed with glyphosate products (such as Touchdown or Roundup brands) for control of volunteer corn including volunteer glyphosate-resistant corn in glyphosate-tolerant soybeans (including Roundup Ready soybeans). Apply Fusion Herbicide at 4-6 fl. oz./A. Use the 4 fl. oz./A rate only under the following conditions:

- Application under favorable soil moisture and humidity conditions which normally occur within a few days after rainfall or irrigation.
- Avoid extreme air temperatures.
- Application when volunteer corn is less than 12 inches tall.
- Application with 0.25% v/v crop oil concentrate (COC). This COC is in addition to the additives required by the glyphosate product.

Use the higher rate if the above conditions do not exist.

Application of this tank mix on soybean varieties that have not been genetically modified to tolerate glyphosate-based products will result in severe injury or plant death.

## TANK MIX AND SEQUENTIAL APPLICATIONS FOR COTTON

Tank mixes with other herbicides labeled for use in cotton are not recommended unless specified on this label or other supplemental labeling.

### Glyphosate-Tolerant Cotton (including Roundup Ready Cotton) Tank Mix

Fusion Herbicide can be tank mixed with glyphosate products (such as Touchdown or Roundup brands) for control of volunteer corn including volunteer glyphosate-resistant corn in glyphosate-tolerant cotton (including Roundup Ready cotton). Apply Fusion Herbicide at 4-6 fl. oz./A. Use the 4 fl. oz./A rate only under the following conditions:

- Application under favorable soil moisture and humidity conditions which normally occur within a few days after rainfall or irrigation.
- Avoid extreme air temperatures.
- Application when volunteer corn is less than 12 inches tall.
- Application with 0.25% v/v crop oil concentrate (COC). This COC is in addition to the additives required by the glyphosate product.

Use the higher rate if the above conditions do not exist.

Application of this tank mix on cotton varieties that have not been genetically modified to tolerate glyphosate based products will result in severe injury or plant death.

### Sequential Applications

Fusion Herbicide can be used sequentially with other labeled cotton herbicides. For sequential applications, allow 2 to 3 days after the application of Fusion Herbicide before applying a broadleaf herbicide or mixture. In situations where the broadleaf herbicide or mixture is applied first, apply Fusion Herbicide when the grass weeds begin to develop new leaves (generally around 7 days).

### Cotton Tank Mix and Sequential Use Precautions

- Always read and follow the directions, restrictions and limitations for all products whether used alone, sequentially or in a tank mix. The most restrictive labeling of any product used applies.

## APPENDIX

Scientific names are listed for those weeds referred to in the Fusion Herbicide label.

COMMON NAME	SCIENTIFIC NAME
Barnyardgrass	<i>Echinochloa crus-galli</i>
Broadleaf signalgrass	<i>Brachiaria platyphylla</i>
Brome, Downy	<i>Bromus tectorum</i>
Crabgrass, Large	<i>Digitaria sanguinalis</i>
Crabgrass, Smooth	<i>Digitaria ischaemum</i>
Crabgrass, Southern	<i>Digitaria ciliaris</i>
Crabgrass, Tropical	<i>Digitaria bicornis</i>
Fall Panicum	<i>Panicum dichotomiflorum</i>
Field Sandbur	<i>Cenchrus incertus</i>
Foxtail, Giant	<i>Setaria faberi</i>

COMMON NAME	SCIENTIFIC NAME
Foxtail, Green	<i>Setaria viridis</i>
Foxtail, Yellow	<i>Setaria pumila</i>
Goosegrass	<i>Eleusine indica</i>
Itchgrass	<i>Rottboellia exaltata</i>
Johnsongrass, Seedling	<i>Sorghum halapense</i>
Junglerice	<i>Echinochloa colonum</i>
Quackgrass	<i>Elymus repens</i>
Red Rice	<i>Oryza sativa</i>
Shattercane	<i>Sorghum bicolor</i>
Sorghum Almum	<i>Sorghum almum</i>
Southern Sandbur	<i>Cenchrus echinatus</i>
Southwestern cupgrass	<i>Eriochloa gracilis</i>
Texas Panicum	<i>Panicum texanum</i>
Volunteer Cereals	
V. Barley	<i>Hordeum vulgare</i>
V. Corn	<i>Zea mays</i>
V. Milo	<i>Sorghum bicolor</i>
V. Oats	<i>Avena sativa</i>
V. Rye	<i>Secale cereale</i>
V. Wheat	<i>Triticum aestivum</i>
Wild Oats	<i>Avena fatua</i>
Wild Proso Millet	<i>Panicum miliaceum</i>
Wirestem Muhly	<i>Muhlenbergia frondosa</i>
Witchgrass	<i>Panicum capillare</i>
Woolly cupgrass	<i>Eriochloa villosa</i>

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

### Pesticide Storage

Store in original container only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with sand, earth or synthetic absorbent. Remove to chemical waste area.

### **Pesticide Disposal**

Open dumping is prohibited. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

### **Container Handling [less than 5 gallons]**

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple-rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### **Container Handling [Bulk/Mini-bulk]**

Refillable container. Refill this container with pesticide only. Do not use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.**

---

Boundary®, Fusion®, Flexstar®, Flexstar® GT, Gramoxone Inteon®, Prefix®, Reflex®, Touchdown®, the ALLIANCE FRAME  the SYNGENTA Logo and the PURPOSE ICON  are Trademarks of a Syngenta Group Company.

Basagran®, Prowl®, Pursuit®, Scepter® and Storm® trademarks of BASF Corporation.

FirstRate® and Garlon® trademarks of Dow Agro Sciences.

Canopy®, Classic®, Escort®, Harmony®, Synchrony®, Telar® and Viton® trademarks of E. I. duPont de Nemours & Co., Inc.

Cobra® trademark of Valent USA Corporation.

Ignite® and LibertyLink® trademarks of Bayer CropScience

IMI-Corn® trademark of American Cyanamid Company

Roundup® and Roundup Ready® trademarks of Monsanto Company.

Ultra Blazer® trademark of United Phosphorus, Inc.

©2011 Syngenta

For non-emergency (e.g., current product information), call  
Syngenta Crop Protection at 1-800-334-9481

Manufactured for:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina 27419-8300

**SCP 1059A-L1D 1210  
334497**

GROUP 1 HERBICIDE



# Herbicide

Postemergence Herbicide for Control of Perennial and Annual Grass Weeds

**Active Ingredients:**

Fluazifop-P-butyl	
Butyl(R)-2-[4-[[5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]propanoate	24.15%
Fenoxaprop-P-ethyl (+)-ethyl-2-[4-[6-(chloro-2-benzoxazolyl)oxy]phenoxy]propanoate	6.76%
Other Ingredients*	69.09%
<b>Total:</b>	<b>100.00%</b>

Contains 2 lbs. (+) isomer (fluazifop-P-butyl) and 0.56 lbs. fenoxaprop-P-ethyl active ingredient per gallon.

\*Contains aromatic petroleum distillates.

EPA Reg. No. 100-1059  
EPA Est. 11773-IA-01

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Fusion® and the Syngenta logo are trademarks of a Syngenta Group Company.

©2011 Syngenta

Manufactured for:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina 27419-8300

**SCP 1059A-L1D 1210  
334497**

## 2.5 gallons

Net Contents

## KEEP OUT OF REACH OF CHILDREN.

## WARNING/ AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

### Precautionary Statements

#### Hazards to Humans and Domestic Animals WARNING/AVISO

**WARNING.** Causes skin irritation. Harmful if absorbed through skin or swallowed. Causes moderate eye irritation. Do not get on skin or on clothing. Avoid contact with eyes. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

#### FIRST AID

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **If swallowed:** Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person. **If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If inhaled:** Move person to fresh air. If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

**NOTE TO PHYSICIAN:** Contains petroleum distillate – vomiting may cause aspiration pneumonia.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**HOT LINE NUMBER:** For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident) Call **1-800-888-8372**.

**Environmental Hazards:** This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Do not apply when weather conditions favor drift from target area.

**Ground Water Advisory:** Fluazifop-P-butyl is known to leach through soil into ground water under certain conditions as a result of label use.

This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water.

This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of fluazifop-p-butyl from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

**Physical and Chemical Hazards: Combustible.** Do not use or store near heat or open flame.

**Chemigation:** Do not apply Fusion Herbicide through any type of irrigation system.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

**Pesticide Storage:** Store in original container only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with sand, earth or synthetic absorbent. Remove to chemical waste area.

**Pesticide Disposal:** Open dumping is prohibited. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**Container Handling [less than 5 gallons]:** Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple-rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.



BAR CODE # IS  
(01) 0 07 02941 73290  
LAST DIGIT IS CHECK DIGIT  
UCC/EAN 128