

Veteran[®] 720

Herbicide

Water soluble industrial herbicide for control of brush and broadleaf weeds on rights-of-way, forest brush, fence rows, non-irrigation ditchbanks, wasteland, general farmstead, and similar noncropland areas. Also for pastures and rangelands.

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-Dichlorophenoxyacetic Acid* 24.58%
Dimethylamine salt of Dicamba (3,6-Dichloro-o-Anisic Acid)** 12.82%

OTHER INGREDIENTS: 62.60%
TOTAL: 100.00%

BY ISOMER SPECIFIC METHOD, EQUIVALENT TO:

*2,4-Dichlorophenoxyacetic Acid 20.4%, 1.9 lbs./gal.
**3,6-Dichloro-o-anisic Acid 10.6%, 1.0 lbs./gal.

KEEP OUT OF REACH OF CHILDREN
DANGER / PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire,
or Exposure, Call CHEMTREC
(800) 424-9300

For Medical Emergencies Only,
Call (877) 325-1840

EPA Reg. No. 228-295

Manufactured for
Nufarm Americas Inc.
11901 S. Austin Avenue
Alsip, IL 60803



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER / PELIGRO**

Corrosive. Causes irreversible eye damage. Harmful if swallowed. Do not get in eyes or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are barrier laminate or Viton. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators, flaggers, and other handlers must wear:

- long-sleeved shirt and long pants
- shoes and socks, plus
- goggles or faceshield
- chemical-resistant gloves (except for applicators using groundboom equipment, pilots, and flaggers)
- chemical-resistant apron when applying postharvest drips or sprays to citrus, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This product has properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For use on plants in non-crop and non-timber areas only. Not for use on trees, crops, or other plants being grown for sale or other commercial use, or for commercial seed production, or for the production of timber or wood products, or for research purposes.

This product is a water soluble industrial herbicide for brush and broadleaf weed control, is recommended for use on rights-of-way (utility, railroad, highway, pipeline), nonselective forest brush control, fence rows, non-irrigation ditchbanks, wasteland, general farmstead, and similar non-cropland areas. Also for pastures and rangeland.

Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls worn over short-sleeve shirt and short pants, chemical-resistant footwear plus socks, chemical-resistant gloves made of any water-proof material, chemical-resistant headgear for overhead exposure and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

SENSITIVE CROP PRECAUTIONS

This product may cause injury to desirable trees and plants, particularly beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes and other broadleaf plants when contacting their roots, stems or foliage. These plants are most sensitive to this product during their development or growing stage.

Do not treat areas where either possible downward movement into the soil or surface washing may cause contact of this product with the roots of desirable plants such as trees and shrubs.

To avoid injury to desirable plants, equipment used to apply this product should be thoroughly cleaned (see PROCEDURE FOR CLEANING SPRAY EQUIPMENT on back panel) before reusing to apply any other chemicals.

MIXING AND APPLICATION

This product is a water-soluble formulation which can be applied using water or oil/water emulsions as the carrier. Herbicide adjuvants or other spray additives (emulsifiers, surfactants, wetting agents, drift control agents, or penetrants) such as Redi-Vert™, Bivert®, Cayuse™, R-11®, and R-900®, may be used for wetting, penetration, or drift control. Read and follow all use recommendations and precautions on the label(s) of product(s) used. A COMPATIBILITY TEST (see other panel) should be made prior to tank mixing.

This product may be applied with ground or aerial equipment for broadcast treatments, or with handgun, backpack, or similar types of equipment for treatment of individual clumps or small areas (spot treatments).

NOTE: To prepare oil/water emulsions, shake well before using, half fill spray tank with water. Add a suitable emulsifier. Then, with constant agitation, slowly add the appropriate amount of herbicide followed by the necessary oil (or slowly add a premix of emulsifier and oil). Finally, fill the rest of the tank with water. Maintain vigorous agitation during spray operation to prevent oil and water from forming separate layers.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Additional requirements for aerial applications:

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

Additional requirements for ground boom applications:

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

SPECIES CONTROLLED

When used as directed, this product will control or suppress many herbaceous broadleaf weeds (annual, biennial, and perennial) as well as many unwanted woody plant and vine species. Species controlled include:

ANNUALS

Buckwheat, wild
Carpetweed
Chickweed
Clover
Cocklebur
Daisy, English

Henbit
Knaw
Lambsquarter
Morningglory
Mustard
Pigweed

Purslane
Ragweed
Smartweed
Velvetleaf

BIENNIALS

Ragwort, Tansy

Thistle, Musk

PERENNIALS

Bindweed, Field
Carrot, Wild
(Queen Anne's Lace)
Dock, Curly

Dogfennel
Knapweed, Russian
Milkweed
Ragweed, Perennial

Sorrel, Sheep
Spurge, Leafy
Thistle, Canada
Toadflax, Dalmatian

WOODY BRUSH AND VINES

Alder	Elm	Plum, Wild*	Serviceberry
Ash	Gum	Poplar	Snowberry
Aspen	Hawthorn*	Puncturevine	Spruce
Basswood	Hemlock	Raspberry	Sumac
Beech	Honeysuckle	Redcedar, Eastern*	Sycamore
Birch	Ivy, Poison	Redvine	Trumpet creeper
Blackberry*	Kudzu	Rose, Multiflora*	Waxmyrtle
Cherry	Locust	Sagebrush	Willow
Creeper, Virginia	Maple	Sassafras	Witchhazel
Creosotebush*	Oak	Schinus	Yaupon*
Cucumber tree	Olive, Russian	(Florida Holly,	
Dogwood*	Persimmon	Brazil Peppertree,	
Elderberry	Pine	Christmas-berry)	

*Suppression

RATES

Regardless of the species to be controlled, spray volumes should be high enough to allow for good spray coverage. Make applications when weeds and brush are actively growing. The addition of surfactants can increase control. Biennials are best controlled when treated in the rosette stage. Regrowth may occur on resistant species. Retreatments may be made as needed; however, do not exceed a total of 2 gallons (2 pounds a.i. dicamba + 3.8 pounds a.i. 2,4-D) of this product per treated acre during a growing season. To control additional weed species, this product may be tank mixed with any of the products listed on this label.

HERBACEOUS BROADLEAF WEED CONTROL

Apply 1 to 4 quarts of this product in 20 to 100 gallons of water per treated acre, (0.75 to 3 ounces per 1,000 square feet). When using low-volume application equipment, 3 to 20 gallons of water per acre is acceptable. 1 to 2 quarts (0.75 to 1.5 ounces) of this product is recommended for annuals, 2 to 4 quarts (1.5 to 3.0 ounces) for biennials and easy-to-kill perennials, and 4 quarts for established perennials.

BRUSH AND VINE CONTROL

High Volume Foliar Spot Applications: Mix 1 gallon of this product in enough water to make 100 gallons of spray mix. When using low-volume application equipment, 3 to 20 gallons of water per acre is acceptable. Spray volume applied will depend on the size and density of the brush to be treated. Direct the spray to treat all foliage, stems, and root collars to wet.

Broadcast Applications with Ground Equipment: Apply 1 gallon of this product in 20 to 100 gallons of water per treated acre. When using low-volume application equipment, 3 to 20 gallons of water per acre is acceptable. Spray volume applied will depend on the size and density of the brush to be treated. Spray all foliage, stems and root collars to wet.

Limitations:

- For postemergence annual and perennial weeds in non-cropland, do not apply more than 1 gallon of product per acre per application. Do not apply more than 2 applications per year with a minimum retreatment interval of 30 days. For postemergence woody plants, do not apply more than 1 gallon of product per acre per application. Do not apply more than 1 application per year.
- For postemergence use in pasture and rangeland, do not apply more than 1 gallon of product per acre per application. Do not apply more than 2 applications per year with a minimum retreatment interval of 30 days. If grass is to be cut for hay, Agricultural Use Requirements for the Worker Protection Standard are applicable. Do not cut forage for hay within 7 days of application.

AERIAL APPLICATIONS

Aerial applications may be made to control either herbaceous or woody plants. Apply 1 to 4 quarts of this product (for herbaceous weeds) or 1 gallon of this product (for woody brush and vines) in 5 to 40 gallons of water per treated acre. Coverage is important, so increase spray volume when treating dense stands of brush or weeds.

TANK MIX TREATMENTS

READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, AND OTHER RESTRICTIONS. For broader spectrum control, this product may be tank mixed with one or more of the following herbicides for noncropland uses (e.g. railroad, highway, pipeline, etc.) including forest management, pastures and rangeland applications, if permitted by product labeling (e.g. 2,4-D). Add water to the spray tank prior to the addition of the tank mix products. Do not premix concentrates.

HERBICIDE	RATE (LBS. A.I./TREATED ACRE)
Amitrol*	1.8 to 8.0
Asulam	2.92 to 6.68
Atratal	4.77 to 39.96
Bromacil	1.6 to 24.0
Clorflurecol	2.0 to 3.0
Chlorsulfuron	0.1875 to 2.25 oz.***
Dalapon	4.25 to 12.75
Dicamba	0.25 to 1.75****
Diquat	0.5 to 1.0
Diuron	4.0 to 48.0
Fenac	4.5 to 18.0
Fosamine ammonium	6.0 to 12.0
Glyphosate	0.75 to 3.75
Hexazinone	0.675 to 10.8
Imazapyr**	0.5 to 1.5
Limit®***	0.625
Maleic hydrazide	2.25 to 4.5
Mefluidide	0.25 to 1.0
Metsulfuron Methyl ¹	0.3 to 0.9***
MSMA	1.0 to 2.475
Paraquat*	0.5
Picloram*	2.0 to 3.0
Simazine*	4.8 to 40.0
Sulfometuron methyl	0.75 to 9.0 oz.***
Tebuthiuron	1.0 to 16.0
Triclopyr	0.75 to 12.0
2,4-D	0.475 to 3.5*****
2,4-DP	0.5 to 11.1

Due to variations that may occur in formulated products and specific use ingredients (e.g. water supplies) a COMPATIBILITY TEST as described below is recommended prior to actual tank mixing.

* Restricted use pesticides limited to certified applicators.

** Limit does not have a common name.

*** Chlorsulfuron = 0.1875 to 2.25 ounces a.i./acre (0.25 to 3.0 ounces product/acre)

Sulfometuron methyl = 0.75 to 9.0 oz. a.i./acre (1.0 to 12.0 ounces product/acre)

Metsulfuron methyl = 0.3 to 0.9 oz. a.i. (0.5 to 1.5 ounces product/acre)

**** Do not exceed a total of 1 pound a.i. dicamba per treated acre per growing season. (This product contains 1 pound. a.i. dicamba per gallon.)

***** Use of extremely hard water (500 ppm) may cause this product to form a precipitate when used in a tank mix with this product. Do not exceed 4 pounds total 2,4-D acid equivalent per acre per application per site. Do not exceed 16 pounds total 2,4-D acid equivalent per treated acre per growing season. (This product contains the equivalent to 1.9 pounds per gallon 2,4-D acid.)

¹Using this product and Metsulfuron Methyl to reduce the development and spread of resistant biotypes and problem weeds such as Kochia and Russian thistle: Some commonly resistant weeds typically require multiple spray applications to obtain adequate control. To reduce the number of applications required, applicators can utilize a mixture of Metsulfuron Methyl and this product. Apply as a tank mix for postemergence as follows: Add 0.5 ounce of Metsulfuron Methyl to 1.5 quarts of this product per acre. A non-ionic or silicone surfactant may be used for wetting, penetration or drift control.

NOTE: All intended tank mix combinations should be used only in recommended areas on the same broadleaf weed species found on both labels. For application methods and other use specifications, use the most restricted limitations from labeling of both products.

COMPATIBILITY TEST

Before mixing in the spray tank, it is advisable to test compatibility by mixing all components in a small container in proportionate quantities (see following table).

AMOUNT OF HERBICIDE TO ADD TO ONE PINT OF SPRAY CARRIER (ASSUMING VOLUME IS 25 GALLONS PER ACRE)

<u>HERBICIDE FORMULATIONS</u>	<u>RATE PER ACRE</u>	<u>LEVEL TEASPOONS</u>
Dry	1 pound	1-1/2
Liquid	1 pint	1/2

If herbicide(s) do not ball-up or form flakes, sludge, gels, oily films or layers, or other precipitates, then the tested components are compatible. Usually incompatibility in any of the above described forms will occur within 5 minutes after mixing.

If components are incompatible, the use of a compatibility agent is recommended. Rerun the above COMPATIBILITY TEST with a suitable compatibility agent (1/4 teaspoon is equivalent to 2 pints per 100 gallons of spray solution).

PROCEDURE FOR CLEANING SPRAY EQUIPMENT

The steps listed below are suggested for thorough cleaning of spray equipment following applications of this product or tank mixes of this product plus 2,4-D Amine.

- 1) Hose down thoroughly the inside as well as outside surfaces of equipment while filling the spray tank half full of water. Flush by operating sprayer until the system is purged of the rinse water.
- 2) Fill tank with water while adding 1 quart of household ammonia or 1/4 pint of Neutral-Clean™ for every 25 gallons of water. Operate the pump to circulate the ammonia solution through the sprayer system for 15 to 20 minutes and discharge a small amount of the ammonia solution through the boom and nozzles. Let the solution stand for several hours, preferable overnight.
- 3) Flush the solution out of the spray tank through the boom.
- 4) Remove the nozzles and screens and flush the system with two full tanks of water.

The steps listed below are suggested for thorough cleaning of spray equipment used to apply this product as a tank mix with wettable powders (WP), emulsifiable concentrates (EC), or other types of water-dispersible formulations. Tank mixing this product with water-dispersible formulations, requires the use of a water/detergent rinse.

- 5) Complete step 1.
- 6) Fill tank with water while adding 2 pounds of detergent for every 40 gallons of water. Operate the pump to circulate the detergent solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
- 7) Flush the detergent solution out of the spray tank through the boom.
- 8) Repeat step 1, and follow with steps 2, 3 and 4.

Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et. al. v. EPA, C01-0132C, (W.D. WA). For further information, please refer to <http://www.epa.gov/espp/>.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always store pesticides in original container and in a secured warehouse or storage building. Do not store near open containers of fertilizer, seed or other pesticides. Store at temperature above 32°F. If allowed to freeze, rewarm to 40°F, remix thoroughly before using. This does not alter this product. Containers should be opened in well-ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law and may contaminate ground water. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

NOTE: This product is available in multiple containers. Refer to the Net Contents section of this products labeling for the applicable "Nonrefillable" or "Refillable" designation. Follow the container disposal instructions below that apply to your container type / size.

Nonrefillable Containers 5 Gallons or Less: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

STORAGE AND DISPOSAL *(continued)*

Nonrefillable containers larger than 5 gallons: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. If recycling or reconditioning is not available, puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers larger than 5 gallons: Refillable container: Refill this container with pesticide only. Do not reuse this container for any other purpose. Close all openings and replace all caps. Contact Nufarm's Customer Service Department at 1-800-345-3330 to arrange for return of the empty refillable container. If not returned to Nufarm, triple rinse the empty container and offer it for recycling if available. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Local conditions may affect the uses of this chemical as shown on this label. Consult State Experiment Station or Extension Service weed specialist for specific recommendations for local weed problems and for information on possible lower dosages.

WARRANTY DISCLAIMER

The Directions for Use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV110211)

Veteran is a registered trademark of Nufarm Americas Inc.
All other trademarks are the property of their respective owners.